




An introduction to high energy heavy ion physics

COSMOLOGY MARCHES ON




1 small bang in the ALICE experiment




ALICE

Pb+Pb @ $\sqrt{s} = 2.76$ ATeV
2010-11-08 11:30:46
Fill : 1482
Run : 137124
Event : 0x00000000D3BBE693



Matter particles

All ordinary particles belong to this group

These particles existed just after the Big Bang. Now they are found only in cosmic rays and accelerators

LEPTONS		
FIRST FAMILY	Electron Responsible for electricity and chemical reactions; it has a charge of -1	Electron neutrino Particle with no electric charge, and possibly no mass; billions fly through your body every second
SECOND FAMILY	Muon A heavier relative of the electron; it lives for two-millionths of a second	Muon neutrino Created along with muons when some particles decay
THIRD FAMILY	Tau Heavier still; it is extremely unstable. It was discovered in 1975	Tau neutrino not yet discovered but believed to exist

QUARKS		
Up Has an electric charge of plus two-thirds; protons contain two, neutrons contain one	Down Has an electric charge of minus one-third; protons contain one, neutrons contain two	
Charm A heavier relative of the up; found in 1974	Strange A heavier relative of the down; found in 1964	
Top Heavier still	Bottom Heavier still; measuring bottom quarks is an important test of electroweak theory	

Force particles

These particles transmit the four fundamental forces of nature although gravitons have so far not been discovered

Gluons
Carriers of the strong force between quarks

Felt by: quarks

The explosive release of nuclear energy is the result of the strong force

Photons
Particles that make up light; they carry the electromagnetic force

Felt by: quarks and charged leptons

Electricity, magnetism and chemistry are all the results of electro-magnetic force

Intermediate vector bosons
Carriers of the weak force

Felt by: quarks and leptons

Some forms of radio-activity are the result of the weak force

Gravitons
Carriers of gravity


Felt by: all particles with mass

All the weight we experience is the result of the gravitational force




The focus of this talk is quark and gluon physics

- The strong interaction is very complex!




Quarks and gluons couple strong:

QUESTIONS



Complex vacuum:



CONFINEMENT



Quantum Chromo Dynamics (QCD)

3 color charges (red, green, blue)


Hadrons have to be colorless

Baryons have all 3 colors

Mesons has a color and an anti-color

A single quark cannot be observed because it has color!

The quarks are confined inside the hadrons!







QCD & Confinement

- The strong interaction potential
 - Compare the potential of the strong & e.m. interaction

$$V_{em} = -\frac{c}{r} \quad V_s = -\frac{c'}{r} + kr \quad c, c', k \text{ constants}$$

- Confining term arises due to the self-interaction property of the colour field. $k \sim 1 \text{ GeV/fm}$







Question

- What is “the self-interaction property” of the strong force?



Exercise: How big is k ?

- $k=1\text{GeV}/\text{fm}$
- What force does that correspond to in kilograms?
 - $mg = 1\text{ GeV}/\text{fm} \Rightarrow m=?$






Consequences of 10 ton force!

- This is why QCD is also called the strong interaction
 - QCD can bind together quarks even though they are EM repulsed
- QCD is for low energies non-perturbative
 - We know the theory but we cannot solve it!
 - We don't know how to describe hadronic properties with QCD
- But at high energies (small distances $\ll 1$ fm) we can use perturbative QCD
- Idea: Can we create high energy matter where the quarks and gluons are the fundamental degrees of freedom
 - This is also the phase of matter in the universe around 1 micro second after the big bang!
 - It is first after this time that quarks and gluons “crystallize” into hadrons




Lattice QCD results (Numerical non-perturbative)


QCD energy density



Heavy quark potential



At $T \sim T_c$ the strong potential is screened so e.g. $c+c\text{-bar}$ states can disassociate.





Exercise: What is the high energy limit of QCD?

$$\epsilon_{QCD} = \frac{\pi^2}{30} \left(\begin{matrix} \text{????} \\ \uparrow \\ \text{Bosonic degrees} \\ \text{of freedom (gluons)} \end{matrix} + \frac{7}{8} \begin{matrix} \text{??????} \\ \uparrow \\ \text{Fermionic degrees} \\ \text{of freedom (quarks)} \end{matrix} \right) T^4$$

Bosonic degrees
of freedom (gluons)

Fermionic degrees
of freedom (quarks)




Answer:

$$\epsilon_{QCD} = \frac{\pi^2}{30} \left(2 \times 8 + \frac{7}{8} 2 \times 2 \times 3 \times 3 \right) T^4$$

↑
Gluon spin and color

↑
(Anti+)quark spin, color and flavor




This suggests that the Quark Gluon Plasma should behave as a gas of quarks and gluons!



QCD phase diagram

$T \sim 170 \text{ MeV}$ ($\sim 1 \text{ GeV}/\text{fm}^3$)
 $1 \text{ eV} = 11605 \text{ K}$
 $T \sim 2,000,000,000,000 \text{ K}$
(T core sun:
 $16,000,000 \text{ K}$)




- By colliding heavy ions we hope to create (and study the characteristics of) a new phase of matter called the Quark Gluon Plasma (where quarks and gluons are deconfined)





Heavy ion collisions: The study of high energy QCD

The evolution of a heavy ion collision



- By colliding heavy ions it is possible to create a large ($\gg 1\text{fm}^3$) zone of hot and dense QCD matter
- Goal is to create and study the properties of the Quark Gluon Plasma
- Experimentally only the final state particles are observed, so the conclusions have to be inferred via models

QGP – the phase of the universe 1 micro second after The Big Bang





anti-quark He helium
 e⁻ electron Li lithium

© Fraunhofer



Assumed knowledge


- Accelerators to produce the high energy beams
 - Relativistic Heavy Ion Collider at Brookhaven National Laboratory (outside new York)
 - Large Hadron Collider at CERN (near Geneva)
- Experiments to detect and reconstruct the final state particles
 - PHENIX and STAR at the Relativistic Heavy Ion Collider
 - ATLAS and ALICE at the Large Hadron Collider





Heavy Ion Jargon

Centrality (ex. for Au+Au):




$$\text{Participants} = 2 \cdot 197 - \text{Spectators}$$


- The total energy is proportional to the participant
- The number of parton-parton (quark-quark, quark-gluon, gluon-gluon) is proportional to the binary collisions

Example:



- 6 participant
- 8 binary collisions
- (pp has 2 participant and 1 binary collision)







Peripheral Event

From real-time Level 3 display.


color code \Rightarrow energy loss






Mid-Central Event


From real-time Level 3 display.






Central Event

From real-time Level 3 display.



ALICE first collisions: 8/11-2010

Factor 14 jump in energy!



Pb+Pb @ $\sqrt{s} = 2.76$ ATeV

2010-11-08 11:30:46

Fill : 1482

Run : 137124

Event : 0x00000000D3BBE693






What happens when we collide *pp* and *Pb-Pb*


2 answers!

SOFT




Non-perturbative physics
(know the equations but not how to solve them)
Bulk properties (=medium)

HARD





Perturbative physics
(theoretical predictions)
Rare processes
jets (=probes)






Phenomenological model of soft physics e.g. Lund string model



a) QED or QCD ($r < 1$ fm)




b) QCD ($r > 1$ fm)






Charged Particle Multiplicity

$dN/d\eta$



According to Bjorken:

$$\epsilon \approx \frac{1}{A_t} \frac{dN}{d\eta} \frac{1}{\tau} \langle E_t \rangle$$




Estimate the energy density, assume $\langle E_t \rangle \sim 0.5 \text{ GeV}$,



“Measured” initial energy density


Bjorkens hydrodynamic formula for thermalized energy density in terms of measured transverse energy E_T

$$\varepsilon_{Bj} = \frac{1}{\pi R^2} \frac{1}{c\tau_0} \left(\frac{dE_T}{dy} \right)$$



PHENIX: Central Au Au yields



$$\left\langle \frac{dE_T}{d\eta} \right\rangle_{\eta=0} = 503 \pm 2 \text{ GeV}$$



Formation(thermalization) time ?



LHC first results from ALICE



- Multiplicity at *mid-rapidity* increases a factor ~ 2 from RHIC to LHC (energy density a factor ~ 2) but most likely also widens significantly
- Centrality dependence is similar as at lower energy



Break


- Please take 1 minute to write down on a piece of paper what the muddiest point so far has been





Short recap




- We want to prove that the matter formed in heavy ion collisions is the expected Quark Gluon Plasma predicted by theory
 - a high energy gas/phase of quarks and gluons
- Problem: We have to derive this from the final state particles that are emitted after the system has cooled of
- We have shown that the energy density derived from the charged particle density is larger than the energy density required from QCD numerical simulations
 - **Necessary condition, but not sufficient condition**
- We want now to show that the matter formed is strongly interacting and that it shows quark and gluon degrees of freedom



Elliptic flow (v_2)


unique in heavy ion collisions

Fourier decomposition:
 $dN/d\phi = 1 + 2 V_2 \cos(2 \Delta\phi)$







Example from atomic physics: Releasing Lithium atoms from a trap



<http://www.physics.ncsu.edu/jet/index.html>



Elliptic flow exercise(s)





- Why is the elliptic flow sensitive to early interactions after the hot and dense matter has been formed?
 - Hint: How would the phi distribution look if there were no interactions
- Bonus question: Why is the flow generated in the event plane and not transverse to that?
 - Hint: How is the matter density distributed in the overlap region



Answers


- Each nucleon-nucleon interaction produces on average a spherical symmetric distribution. Only by interacting elliptic flow is generated

Zhang, Gyulassy, Ko,
Phys. Lett. B455 (1999) 45




- Flow is strongest in the event plane because of the stronger matter gradient – hydrodynamic explanation

SCIENCE Vol: 298 2179 (2002)





Elliptic flow at RHIC is “Maximal”



- Relativistic hydrodynamic predicts elliptic flow
 - The high energy medium interacts very strongly immediately after being formed
 - Medium does not behave as a gas, but an almost perfect fluid!



The QGP is less like a crowd and more like a synchro team





■ Big theoretical challenge:

- how to go from initial random collisions to organized state in a VERY short time ($<1\text{fm}/c \sim 10^{-23}\text{s}$)
- Remains to be understood





LHC first results from ALICE



- *Differential flow at LHC is very similar to RHIC!*
 - Hydrodynamic limit?
- The total magnitude is larger because system is harder!
- Question: Where is QCD dynamics?




Hard probes (pQCD): parton-parton interactions






Heavy Ion Jargon Revisited

Centrality (ex. for Au+Au):





$$\text{Participants} = 2 \cdot 197 - \text{Spectators}$$

- parton-parton collisions are proportional to binary collisions
- Exercise: Why is the number of binary collisions in central collisions proportional to $A^{4/3}$ while the number of participants is A ?
 - Hint: What is the average amount of nuclear matter covered in the “target” nuclei?








The nuclear modification factor for pions (1/2)



$$R_{AA} = \frac{d^2 N^{AA} / d p_T dy}{\langle N_{bin} \rangle d^2 N^{NN} / d p_T dy}$$





Nbin






The nuclear modification factor for pions (2/2)




- In central collisions we observe only 20% of the remnants from parton-parton collisions that we expected to observe!
- What happens to the rest?
 - They lose energy as they go through the high energy matter!
 - This is the QCD signature we looked for!
- But first let us consider other alternatives!



Could the binary scaling be wrong?





Source of direct photons



- Direct photons does not interact with final state hadronic matter!
- Direct photons shows no nuclear modification and therefore confirm binary scaling of hard processes!





New “standard candle” at LHC: ATLAS measures Z bosons



- The Z does not interact strongly and so can also be used to check binary scaling at LHC







Could it be an initial state effects? Au+Au vs d+Au








Could it be an initial state effects? Au+Au vs d+Au








Could it be an initial state effects? Au+Au vs d+Au





Could it be an initial state effects? Au+Au vs d+Au




So it must be a final state effect!




The suppression is due to energy loss in the medium

Side view




Most jets are created back to back!

Back view



Adler *et al.*, PRL90:082302 (2003), STAR




$4 < p_T(\text{trig}) < 6 \text{ GeV}/c$

$p_T(\text{assoc}) > 2 \text{ GeV}/c$





The suppression is due to energy loss in the medium

Side view



Most jets are created back to back!

Back view



$4 < p_T(\text{trig}) < 6 \text{ GeV}/c$



$p_T(\text{assoc}) > 2 \text{ GeV}/c$

A large energy loss requires a QCD interacting medium, i.e., a colored medium!





Au+Au vs d+Au

Hot vs cold nuclear matter





All 4 experiments published together in PRL:



No suppression seen in d+Au
 → Quarks and gluons loose/radiate energy as they interact with the colored quarks and gluons of the created matter.
 This suggests that the quark gluon plasma has been discovered!




LHC first results from ALICE



- The nuclear modification factor is lower than at RHIC suggesting that the energy loss is larger.
- The rise with p_T was not observed at RHIC and could give new insight into the energy loss mechanism



New realm at LHC: ATLAS measures very high p_T jets



- Jet asymmetry confirms picture from RHIC – away side jet is absorbed/modified by the medium
 - Advantage of jets is that they “map” onto the QCD degrees of freedom: quarks and gluons




Summary:


- Hard experimental work at RHIC has led to the conclusion that a Quark Gluon Plasma is most likely produced in central collisions of gold on gold! BUT
- Theoretical models are not very constrained by the data as they use many phenomenological inputs
 - New excitement: Can string theory describe non-perturbative QCD?
- Many observations suggest that the picture is more complicated (Quark Gluon Plasma is not like we expected)
 - Particularly heavy quark data challenges many models
- A lot of new results from LHC has to be digested by the community. This will hopefully help constrain the models.



RAA for identified charged hadrons (Lund)



Pushing the separation to the relativistic rise




Baryon anomaly in central PbPb. Quark recombination?




Baryon anomaly not observed at high p_T as speculated pre-LHC.

$p_T \rightarrow$

pp





central PbPb






RAA identified hadrons





- Identified spectra from $2-3 < p_T < 20$ GeV/c from Lund analysis. First time presented by Antonio Ortiz Velasquez (Lund) at Quark Matter 2012, August 12-18, Washington.
- Extends ALICE unique PID capabilities to the hard regime




Heavy quark thermometer



Observation of sequential suppression of Υ ($b+b$ -bar) family



Note: $6.5 < p_T < 30$ GeV for J/ψ and $\psi(2s)$



Expected in terms of binding energy

Unfortunately heavy quark results are more complex when systematically studied!